



EpiPen AAI Instructions

ANAPHYLAXIS

HOW TO USE EPIPEN AAIS

If you think someone is have an anaphylactic reaction,
give the AAI without delay. It will not harm them.

Always consider anaphylaxis in a food-allergic person even if there are no signs of a rash, hives or swelling.

1. Remove the blue safety cap

Grasp the EpiPen in your dominant hand and remove the blue safety cap by pulling straight up. Remember: Blue to the Sky, Orange to the Thigh!



2. Position the orange tip

Hold the EpiPen at 90°, approximately 10cm away from the leg, with the orange tip pointing towards the outer thigh.

3. Administer the EpiPen AAI

Jab the EpiPen firmly into the outer thigh at a right angle. Hold firmly for 3 seconds, before removing and safely discarding.



4. Once the EpiPen AAI has been administered call 999

Ask for an ambulance and say "ana-fill-axis".

5. Lie the person down with legs raised immediately

If the person is not already lying down, they should do so, with legs raised if possible.

If breathing is difficult, allow them to sit. If they have vomited or feel sick, gently turn them on their side.



6. If there are no signs of improvement after 5 minutes, use a second EpiPen AAI

The person should remain still and lying down until the ambulance arrives. Don't try to get up, even if you start to feel better.

7. Start CPR

If there are no signs of life, start CPR immediately until help arrives.



For more information
on EpiPen AAIs >>



Sign up to the free expiry alert service
and receive reminders by text or email when
your EpiPen is about to expire >>

